



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08196 - Goat Anti-Proenkephalin (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: proenkephalin

Official Symbol: PENK

Accession Number(s): NP_001002927.1

Non-Human GeneID(s): 18619 (mouse), 29237 (rat)

Immunogen

Peptide with sequence C-YKDSSKQDESH, from the internal region of the protein sequence according to NP_001002927.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Preliminary experiments gave an approx 40kDa band in Rat Brain lysates after 0.2µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 30.8kDa according to Mouse NP_001002927.1 and 30.9kDa according to Rat NP_058835.1. The 40kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

IHC: Frozen section of Rat Brain (Striatum) shows staining of dense enkephalinergic axon plexus. Frozen section of the Mouse Brain (Stria terminalis) shows staining of a dense enkephalinergic fibre network in the bed nucleus. Recommended concentration, 0.02-0.05µg/ml.

Species Reactivity

Tested: Mouse, Rat

Expected from sequence similarity: Mouse, Rat

EB08196 (0.02µg/ml) staining of PFA-perfused cryosection of Rat Striatum. HRP-staining with Ni-DAB after Biotin-SP-anti-goat amplification. Data obtained by Prof. Erik Hrabovszky, Inst, Exp, Med., Budapest, Hungary.

EB08196 (0.02µg/ml) staining of PFA-perfused cryosection of Mouse Stria terminalis. HRP-staining with Ni-DAB after Biotin-SP-anti-goat amplification. Data obtained by Prof. Erik Hrabovszky, Inst, Exp, Med., Budapest, Hungary.