

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09551 - Goat Anti-RAP80 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: UIMC1, ubiquitin interaction motif containing 1, RAP80, X2HRIP110, OTTHUMP00000161441, receptor associated protein 80, retinoid x receptor interacting protein

Official Symbol: UIMC1

Accession Number(s): NP_057374.3

Human GeneID(s): [51720](#)

Non-Human GeneID(s): 20184 (mouse), 290997 (rat)

Immunogen

Peptide with sequence C-TRRQKEAKTKSDSG, from the internal region of the protein sequence according to NP_057374.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments gave an approx 60kDa band in Human Heart and Spleen lysates and in lysates of cell lines A431, HeLa and HepG2 after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 79.7kDa according to NP_057374.3. The 60kDa band was successfully blocked by incubation with the immunizing peptide.

IHC: Paraffin embedded Human Testis. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig



EB09551 (3.75µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.