

#### **UK Office**

**Everest Biotech Ltd** 

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05744 - Goat Anti-SART3 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** SART3, squamous cell carcinoma antigen recognised by T cells 3, TIP110, KIAA0156, p110(nrb), RP11-13G14, tat-interacting protein of 110 kDa, DSAP1,

MGC138188, P100, p110 Official Symbol: SART3

Accession Number(s): NP\_055521.1

Human GeneID(s): 9733

#### **Immunogen**

Peptide with sequence C-MSNADFAKLFLRK, from the C Terminus of the protein sequence according to NP\_055521.1.

Please note the <u>peptide</u> is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 170kDa band observed in lysates of cell lines Daudi, Jurkat and MOLT4 (calculated MW of 111kDa according to NP\_055521.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Yang et al, Cancer Res. 1999 Aug 15;59(16):4056-63; PMID: 10463607). Recommended concentration: 0.01-0.03μg/ml.

- -

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB05744 staining (0.01μg/ml) of Jurkat cell lysate (RIPA buffer, 30μg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.