

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08932 - Goat Anti-SEPT6 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SEPT6, septin 6, KIAA0128, MGC16619, MGC20339, SEP2, SEPT2, septin 2

Official Symbol: Sep-06

Accession Number(s): NP_665798.1; NP_055944.2; NP_665801.1

Human GeneID(s): [23157](#)

Important Comments: This antibody is expected to recognize the reported isoforms NP_665798.1, NP_055944.2 and NP_665801.1. Reported variants NP_665799.1 and NP_665798.1 represent identical protein.

Immunogen

Peptide with sequence C-DEVNAFKQRKTA, from the internal region of the protein sequence according to NP_665798.1; NP_055944.2; NP_665801.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.


Western blot: Approx 50kDa band observed in Human Testis, Thyroid Gland and MOLT4 cell lysates (calculated MW of 49.7kDa according to NP_055944.2). Recommended concentration: 0.03-1µg/ml.

IHC: In paraffin embedded Human Tonsil shows specific staining in dividing cells just outside a germinal centre. Recommended concentration, 3-5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Cow, Dog



EB08932 (1 μ g/ml) staining of Human Testis lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.



EB08932 (3.8 μ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.