

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09143 - Goat Anti-SLC12A1 / NKCC Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SLC12A1, solute carrier family 12 (sodium/potassium/chloride transporters), member 1, BSC1, MGC48843, NKCC2, NKCC2A variant A, Na-K-2Cl cotransporter, sodium potassium chloride cotransporter 2 Official Symbol: SLC12A1 Accession Number(s): NP_000329.2 Human GeneID(s): <u>6557</u>

Immunogen

Peptide with sequence C-PHYEETSFGDEAQKR, from the internal region of the protein sequence according to NP_000329.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000. **IHC:** In paraffin embedded Human Kidney shows strong membrane staining in some CT cells. Recommended concentration, 3-6μg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human

Specific Reference

This antibody has been successfully used in WB and IF in Mouse:

Alshahrani S, Almutairi MM, Kursan S, Dias-Junior E, Almiahuob MM, Aguilar-Bryan L, Di Fulvio M. Increased Slc12a1 expression in β -cells

and improved glucose disposal in Slc12a2 heterozygous mice. J Endocrinol. 2015 Dec;227(3):153-65.

PMID: 26400961

EB09143 (3.8ug/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.