

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06896 - Goat Anti-SPFH2 / ERLIN2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ERLIN2, SPFH2, ER lipid raft associated 2, SPFH domain family, member 2, HGNC:1356, C8orf2, MGC87072, chromosome 8 open reading frame 2, Erlin-2, MGC87072

Official Symbol: ERLIN2

Accession Number(s): NP_009106.1

Human GeneID(s): [11160](#)

Non-Human GeneID(s): 290823 (rat)

Important Comments: This antibody is expected to recognise isoform 1 (NP_009106) only.

Immunogen

Peptide with sequence C-EDEPLETATKEN, from the C Terminus of the protein sequence according to NP_009106.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 43kDa band observed in Human Breast lysates and in lysates of Human Breast Cancer (calculated MW of 37.8kDa according to NP_009106.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Pearce et al, J Biol Chem. 2007 Jul 13;282(28):20104-15.; PMID: 17502376). Recommended concentration: 0.3-1µg/ml.

IHC: Paraffin embedded Human Breast. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Rat

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa

25kDa
20kDa

15kDa
10kDa

EB06896 staining (1 μ g/ml) of Human Breast lysate (RIPA buffer, 30 μ g total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



EB06896 (3.75 μ g/ml) staining of paraffin embedded Human Breast. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.