

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11284 - Goat Anti-SPON2 (301-315) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: differentially expressed in cancerous and non-cancerous lung cells 1, DIL1, DIL-1, DKFZp686G21139, FLJ16313, FLJ22401, FLJ34460, MINDIN, M-SPONDIN, spondin 2, extracellular matrix protein, spondin-2, SPON2

Official Symbol: SPON2

Accession Number(s): NP_036577.1

Human GeneID(s): [10417](#)

Non-Human GeneID(s): 100689 (mouse), 171569 (rat)

Important Comments: Reported variants represent identical protein: NP_036577.1, NP_001121797.1, NP_001185950.1

Immunogen

Peptide with sequence C-RTRYVRVQPANNGSP, from the internal region (near C Terminus) of the protein sequence according to NP_036577.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.


Western blot: Approx 35kDa band observed in lysates of cell line A549 (calculated MW of 35.8kDa according to NP_036577.1). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Tonsil. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB11284 (1 μ g/ml) staining of A549 lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB11284 (5 μ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.