

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07264 - Goat Anti-TAK1 / MAP3K7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MAP3K7, TAK1, mitogen-activated protein kinase kinase kinase 7, TGF1a, TGF-beta activated kinase 1, transforming growth factor-beta-activated kinase 1

Official Symbol: MAP3K7

Accession Number(s): NP_003179.1; NP_663304.1

Human GeneID(s): [6885](#)

Non-Human GeneID(s): 26409 (mouse)

Important Comments: This antibody is expected to recognise isoform A (NP_003179.1) and isoform B (NP_663304.1).

Immunogen

Peptide with sequence C-AELDQDEKDQQNT, from the internal region of the protein sequence according to NP_003179.1; NP_663304.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: In transfected HEK293 transiently expressing full-length Human MAP3K7 (myc and DYKDDDDK tagged), a band of approx. 75kDa was observed. No bands were observed in mock-transfected HEK293 and the same band was observed using anti-DYKDDDDK tag antibody. Recommended concentration: 1-2µg/ml.

Purchase the +ve control! The transfected lysate used in QC is available for sale. Please email sales@everestbiotech.com to request datasheet, pricing and delivery information.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat

HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human MAP3K7 with DYKDDDDK tag probed with EB07264 (1ug/ml) in Lane A and probed with anti- DYKDDDDK Tag (1/3000) in lane C. Mock-transfected HEK293 probed with EB07264 (1mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.