

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07587-B - Goat Anti-TCF3 / ITF1, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TCF3, transcription factor 3, E2A, E47, ITF1, TCF-3, VDIR, bHLHb21, NOL1-TCF3 fusion, VDR interacting repressor, class B basic helix-loop-helix protein 21, helix-loop-helix protein HE47, immunoglobulin transcription factor 1, kappa-E2-binding factor, negative vitamin D response element-binding protein, transcription factor 3 (E2A immunoglobulin enhancer binding factors E12/E47), transcription factor 3 variant 3, transcription factor ITF-1, vitamin D receptor-interacting repressor

Official Symbol: TCF3

Accession Number(s): NP_003191.1

Human GeneID(s): [6929](#)

Important Comments: This antibody is expected to recognize reported isoform E12 (NP_003191.1) only.

Immunogen

Peptide with sequence C-KAPRARTSPDEDED., from the internal region of the protein sequence according to NP_003191.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 75kDa band observed in lysates of cell line Daudi (calculated MW of 67.6kDa according to NP_003191.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



Biotinylated EB07587 (0.3µg/ml) staining of Daudi lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.