

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11050 - Goat Anti-TNF-R-II / TNFR2 (mouse) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CD120b, p75, p75 TNF receptor, p75TNFR, p80 TNF-alpha receptor, soluble TNFR1B variant 1, TBPII, TNFBR, TNFR1B, TNFR2, TNF-R2, TNF-R75, TNFR80, TNFR-II, TNF-RII, TNF-R-II, tumor necrosis factor beta receptor, tumor necrosis factor binding protein 2, tumor necrosis factor receptor 2, tumor necrosis factor receptor superfamily member 1B, tumor necrosis factor receptor superfamily, member 1B, tumor necrosis factor receptor type II, Tnfrsf1b

**Official Symbol:** Tnfrsf1b

**Accession Number(s):** NP\_035740.2

**Non-Human GeneID(s):** 21938 (mouse), 156767 (rat)

**Important Comments:** The immunizing peptide represents part of a potential extracellular domain.

### Immunogen

Peptide with sequence KHFCNKTSDTVCAD, from the internal region of the protein sequence according to NP\_035740.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 55kDa band observed in Mouse and Rat Spinal Cord lysates (calculated MW of 50.3kDa according to Mouse NP\_035740.2). Recommended concentration: 0.2-0.6µg/ml.

### Species Reactivity

**Tested:** Mouse, Rat

**Expected from sequence similarity:** Mouse, Rat

EB11050 (0.2µg/ml) staining of Rat Spinal Cord lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.