



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB12033-B - Goat Anti-TNNT3, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** TNNT3, troponin T3, fast skeletal type, TNNTF, beta TnTF, beta-TnTF, fTnT, fast skeletal muscle troponin T, troponin T type 3 (skeletal, fast), troponin-T3, skeletal, fast

**Official Symbol:** TNNT3

**Accession Number(s):** NP\_006748.1; NP\_001036246.1; NP\_001036247.1

**Human GeneID(s):** [7140](#)

**Important Comments:** This antibody is expected to recognize isoform 1, 2 and 4 (NP\_006748.1; NP\_001036246.1; NP\_001036247.1). Reported variants represent identical protein: NP\_001284575.1, NP\_001036247.1

### Immunogen

Peptide with sequence CTTLSRIDQAQKHSK., from the C Terminus of the protein sequence according to NP\_006748.1; NP\_001036246.1; NP\_001036247.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Approx 35kDa band observed in Human, Mouse and Rat Skeletal Muscle lysates (calculated MW of 30.6kDa according to NP\_006748.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Marian et al, Circ Res. 1997 Jul;81(1):76-85. PMID: 9201030.). Recommended concentration: 0.01-0.03µg/ml. See non-biotinylated parental product's datasheet for further QC data.

### Species Reactivity

**Tested:** Human, Mouse, Rat

**Expected from sequence similarity:** Human, Mouse, Rat



Biotinylated EB12033 (0.01 $\mu$ g/ml) staining of Human (A), Mouse (B) and Rat (C) Skeletal Muscle lysates (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.