

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05129 - Goat Anti-TRPM7 / LTRPC7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TRPM7, transient receptor potential cation channel, subfamily M, member 7, CHAK, CHAK1, LTRPC7, FLJ20117, TRP-PLIK, LTRPC ion channel family member 7, homolog of mouse transient receptor potential-phospholipase C-interacting kinase CHaK hypothetical protein FLJ20117, FLJ25718

Official Symbol: TRPM7

Accession Number(s): NP_060142.3

Human GeneID(s): [54822](#)

Immunogen

Peptide with sequence C-TKESESTNSVRLML, from the C Terminus of the protein sequence according to NP_060142.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.

IHC: In paraffin embedded Mouse Brain shows neuronal staining in the Hippocampus.
Recommended concentration: 4-6µg/ml.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig



EB05129 (4µg/ml) staining of paraffin embedded Mouse Hippocampus. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining. Similar results were obtained after antigen retrieval at pH9.