

### **UK Office**

**Everest Biotech Ltd** 

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales:

sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08600 - Goat Anti-VGLUT1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: SLC17A7, solute carrier family 17 (sodium-dependent inorganic phosphate cotransporter), member 7, BNPI, VGLUT1, brain-specific Na-dependent inorganic phosphate cotransporter, solute carrier family 17, member 7, vesicular glutamate transporter 1 Official Symbol: SLC17A7 Accession Number(s): NP\_064705.1 Human GeneID(s): <u>57030</u>

### Immunogen

Peptide with sequence C-HDQLAGSDDSEMED, from the internal region of the protein sequence according to NP\_064705.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**IHC:** In paraffin embedded Human Cortex shows vesicular staining amongst the many axons and dendrites. Recommended concentration, 3-5µg/ml. In paraffin embedded Mouse Brain shows staining of the periphery of the ventromedial nucleus in the hypothalamus Recommended concentration, 0.3-1µg/ml.

## **Species Reactivity**

Tested: Human, Mouse Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow EB08600 (3.8µg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB08600 (0.25µg/ml) staining of paraffin embedded Mouse Hypothalamus and Thalamus. Antigen retrieval at 80C for 30min with citrate buffer pH 6, HRP-staining (data kindly provided by Dr. E. Hrabovszky, Budapest, Hungary