

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11598 - Goat Anti-ZNRF1 (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: B830022L21Rik, E3 ubiquitin-protein ligase ZNRF1, MGC101991, nerve injury-induced gene 283 protein, nin283, peripheral nerve injury protein nin283, ring finger protein 42, Rnf42, zinc and ring finger 1, zinc ring finger protein 1, zinc/RING finger protein 1, Znr1, Zrfp1

Official Symbol: Znr1

Accession Number(s): NP_573469.1; NP_001162093.1

Human GeneID(s): [84937](#)

Non-Human GeneID(s): 170737 (mouse), 690769 (rat)

Important Comments: This antibody is expected to recognize reported isoforms a and b (NP_573469.1; NP_001162093.1). Reported variants represent identical protein: NP_001162092.1, NP_573469.1

Immunogen

Peptide with sequence C-SDSTYAHGNGYQET, from the internal region of the protein sequence according to NP_573469.1; NP_001162093.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 28kDa band observed in Mouse embryo and adult Brain lysates (right panel), consistent with the observed band in transfected N2a cells transiently expressing Znr1(left panel) Calculated MW of 23.8kDa according to NP_573469.1.

Recommended concentration: 1-3µg/ml. Data obtained from Dr. Shuji Wakatsuki, National Institute of Neuroscience, Tokyo, Japan

IHC: In paraffin embedded Mouse Cerebral Cortex shows colabeling with Neurofilament M. Recommended concentration, 10µg/ml. Data obtained from Dr. Shuji Wakatsuki, National Institute of Neuroscience, Tokyo, Japan

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

N2a overexpressing Mouse Znr1 (mock transfection in first lane) and probed with EB11598 (1µg/ml), also staining of Mouse Brain lysates (Embryo E14 and adult cerebellum). Primary incubation was 1 hour. Detected by chemiluminescence.

EB11598 (10µg/ml) staining of paraffin embedded Mouse Cerebral Cortex. Microwaved antigen retrieval with citrate buffer pH 6, streptavidine-Alexa 488-staining after biotinylated anti-goat secondary. The Neurofilament M was labeled by Millipore AB1987 (1:100).